

is usually such that all the fuel supply will pass to the engines by way of the main tank, which is refilled as necessary from the auxiliary tanks. In case of emergency, the system selector valve may connect the auxiliary tanks to the engines directly. [F.C.M./J.A.B.]

Fugacity A function introduced by G. N. Lewis to facilitate the application of thermodynamics to real systems. Thus, when fugacities are substituted for partial pressures in the mass action equilibrium constant expression, which applies strictly only to the ideal case, a true equilibrium constant results for real systems as well.

The fugacity f_i of a constituent i of a thermodynamic system is defined by the following equation (where μ_i is the chemical

$$\mu_i = \mu_i^* + RT \ln f_i$$


potential and μ_i^* is a function of temperature only), in combination with the requirement that the fugacity approach the partial pressure as the total pressure of the gas phase approaches zero. At a given temperature, this is possible only for a particular value for μ_i^* , which may be shown to correspond to the chemical potential the constituent would have as the pure gas in the ideal gas state at 1 atm pressure. This definition makes the fugacity identical to the partial pressure in the ideal gas case. For real gases, the ratio of fugacity to partial pressure, called the fugacity coefficient, will be close to unity for moderate temperatures and pressures. At low temperatures and appropriate pressures, it may be as small as 0.2 or less, whereas at high pressures at any temperature it can become very large. See ACTIVITY (THERMODYNAMICS); CHEMICAL EQUILIBRIUM; CHEMICAL THERMODYNAMICS; GAS. [P.J.B.]

Fullerene A molecule containing an even number of carbon atoms arranged in a closed hollow cage. The fullerenes were discovered as a consequence of astrophysically motivated chemical physics experiments that were interpreted by using geodesic architectural concepts. Fullerene chemistry, a new field that appears to hold much promise for materials development and other applied areas, was born from pure fundamental science. See CARBON.

In 1985, fifteen years after it was conceived theoretically, the molecule buckminsterfullerene (C_{60} or fullerene-60) was discovered serendipitously. Fullerene-60 (see illustration) is the archetypal member of the fullerenes, a set of hollow, closed-cage molecules consisting purely of carbon. The fullerenes can be considered, after graphite and diamond, to be the third well-defined allotrope of carbon.

In the fullerene molecule an even number of carbon atoms are arrayed over the surface of a closed hollow cage. Each atom is trigonally linked to its three near neighbors by bonds that delineate a polyhedral network, consisting of 12 pentagons and n hexagons. All 60 atoms in fullerene-60 are equivalent and lie on the surface of a sphere distributed with the symmetry of a truncated icosahedron. The 12 pentagons are isolated and interspersed symmetrically among 20 linked hexagons; that is, the symmetry is that of a modern soccerball. The molecule was named after R. Buckminster Fuller, the inventor of geodesic domes, which conform to the same underlying structural formula. Three of the four valence electrons of each carbon atom are involved in the sp^2 sigma-bonding skeleton, and the fourth p electron is one of 60 involved in a π -delocalized molecular-orbital electron sea that covers the outside (exo) and inside (endo) surface of the molecule. The resulting cloud of π electron density is similar to that which covers the surface of graphite; indeed, the molecule can be considered a round form of graphite. See ELECTRON CONFIGURATION; GRAPHITE.

Fullerene-60 behaves as a soft electrophile, a molecule that readily accepts electrons during a primary reaction step. It can accept three electrons readily and perhaps even more. The molecule can be multiply hydrogenated, methylated, ammonated, and fluorinated. It forms exohedral complexes in which an atom (or group) is attached to the outside of the cage, as well as endohedral complexes in which an atom (for example, lanthanum (La), potassium (K), or calcium (Ca)) is trapped inside the cage.



Structure of C_{60} (buckminsterfullerene).

Fullerene materials have been available for such a short time that applications are yet to be established. However, the properties already discovered suggest that there is likely to be a wide range of areas in which the fullerenes or their derivatives will have uses.

Fullerene-60 was discovered as a direct result of physicochemical investigations that simulated processes occurring in stars and in space. Consequently the likelihood that fullerenes, in particular fullerene-60, and analogs are present in space is a fascinating conjecture. [H.W.K.]

Fuller's earth Any natural earthy material (such as clay materials) which decolorizes mineral or vegetable oils to a sufficient extent to be of economic importance. It has no mineralogic significance. The clay minerals present in fuller's earth may include montmorillonite, attapulgite, and kaolinite.